

This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims:

1. (Withdrawn) A method of making a population of functional hepatocytes for transplantation into a patient, which comprises:
 - a) providing a sample of primary hepatocytes;
 - b) immortalizing the hepatocytes by transforming the hepatocytes with a DNA construct comprising a removable DNA segment containing an oncogene, thereby producing immortalized hepatocytes;
 - c) growing the immortalized hepatocytes; and
 - d) removing the oncogene from the immortalized hepatocytes, the removal resulting in the production of the population of functional hepatocytes for transplantation into the patient.
2. (Withdrawn) The method of claim 1, wherein the hepatocytes are obtained from a human donor.
3. (Withdrawn) The method of claim 1, wherein the oncogene is made removable by flanking it with recombinase target sites, and the removing is accomplished by introducing into the immortalized cells a gene that is expressed to produce a recombinase that specifically recognizes the recombinase target sites.
4. (Withdrawn) The method of claim 3, wherein the recombinase is Cre recombinase and the recombinase target sites are loxP sites.
5. (Withdrawn) The method of claim 1, wherein the oncogene is a gene encoding SV40 large T antigen.
6. (Withdrawn) The method of claim 1, wherein the removable DNA segment further contains a suicide gene, which encodes a gene product that enables destruction of the

immortalized cells by an exogenous agent if the removable DNA segment is not removed from the cells.

7. (Withdrawn) The method of claim 6, wherein the suicide gene is a gene encoding herpes simplex virus thymidine kinase, and the cells are destroyed by exposure to gancyclovir if the removable DNA segment is not removed from the cells.

8. (Currently amended) A population of isolated, immortalized hepatocytes ~~produced by the method of~~ according to claim 16.

9. (Withdrawn) A method of treating a patient for hepatic failure, comprising transplanting into the patient a sufficient quantity of the hepatocytes of claim 8 to provide hepatic function to the patient.

10. (Withdrawn) A method of making a population of functional hepatocytes for supporting hepatic function of a patient, which comprises:

- a) providing a sample of primary hepatocytes;
- b) immortalizing the hepatocytes by transforming the hepatocytes with a DNA construct comprising a removable DNA segment containing an oncogene, a selectable marker gene, and a gene encoding herpes simplex virus thymidine kinase, the genes together being flanked on either side by loxP sites;
- c) growing the immortalized hepatocytes; and
- d) reversing the immortalization of the hepatocytes by removing the DNA segment from the immortalized hepatocytes, the removing being accomplished by introducing into the immortalized hepatocytes a gene encoding Cre recombinase to effect excision of the DNA segment at the loxP sites, the excision resulting in the production of the population of functional hepatocytes for supporting hepatic function of a patient.

11. (Withdrawn) The method of claim 10, wherein the hepatocytes are obtained from a human donor.

12. (Withdrawn) The method of claim 10, wherein the oncogene is a gene encoding SV40 large T antigen.
13. (Withdrawn) The method of claim 10, wherein the selectable marker gene confers hygromycin resistance to cells expressing the gene.
14. (Currently Amended) A population of reverse-immortalized hepatocytes ~~produced by the method of~~ according to claim 25.
15. (Withdrawn) A method of treating a patient for hepatic failure, comprising transplanting into the patient a sufficient quantity of the hepatocytes of claim 14 to provide hepatic function to the patient.
16. (Currently amended) An isolated, immortalized hepatocyte comprising a primary hepatocyte transformed with a DNA construct comprising two recombinase target sites ~~that flank~~ flanking an oncogene which confers immortalization to the hepatocyte, wherein the immortalization is reversible by excision of the oncogene by cleavage at the recombinase target sites when the target sites are exposed to a recombinase that specifically recognizes the target sites and wherein said hepatocyte is entrapped in a matrix disposed to receive cells for transplantation.
17. (Currently amended) The isolated, immortalized hepatocyte of claim 16, wherein the recombinase target sites are loxP sites and the immortalization is reversible by Cre recombinase cleavage at the loxP sites.
18. (Currently amended) The isolated, immortalized hepatocyte of claim 16, wherein the DNA construct further comprises a selectable marker gene.
19. (Currently amended) The isolated, immortalized hepatocyte of claim 16, wherein the DNA construct further comprises a suicide gene, which encodes a gene product

that enables destruction of the immortalized hepatocyte by an exogenous agent if the oncogene is not removed from the cells.

20. (Currently amended) The isolated, immortalized hepatocyte of claim 19, wherein the suicide gene is a gene encoding herpes simplex virus thymidine kinase, and the exogenous agent is gancyclovir.

21. (Currently amended) The isolated, immortalized hepatocyte of claim 16, wherein the hepatocyte is obtained from a human donor.

22. (Currently amended) A cell line comprising a population of the isolated, immortalized hepatocyte of claim 21, which is cell line NKNT-3 wherein said cell line is entrapped in a matrix disposed to receive cells for transplantation.

23. (Withdrawn) The immortalized hepatocyte of claim 20, wherein the hepatocyte is obtained from a rat donor.

24. (Withdrawn) A cell line comprising a population of the immortalized hepatocyte of claim 23, which is cell line C8-B.

25. (Previously Presented) A reverse-immortalized hepatocyte that is functional upon transplantation into a patient, wherein said reverse-immortalized hepatocyte is produced by exposing the DNA construct within the immortalized hepatocyte of claim 16 to a recombinase that excises the oncogene by cleavage at the recombinase target sites and wherein said hepatocyte is entrapped in a matrix disposed to receive cells for transplantation.

26. (Withdrawn) A method of treating a patient for hepatic failure, comprising transplanting into the patient a sufficient quantity of the reverse-immortalized hepatocytes of claim 25 to provide hepatic function to the patient.